

Emergency Tooth Doctor

West	Downtown	East	Vancouver	Hillsboro
14665 SW Millikan Way Beaverton, OR 97006	1505 SW Broadway Portland, OR 97201	12596 SE Stark St Plaza 125, Bldg. N Portland, OR 97233	400 W Fourth Plain Blvd Vancouver, WA 98660	405 SE 8th Ave. Hillsboro, OR 97123
Ph: 503-641-2200 Fax: 503-641-2220	Ph: 503-222-0090 Fax: 503-222-1033	Ph: 503-252-9657 Fax: 503-256-6909	Ph: 360-975-3198 Fax: 360-718-2594	Ph: 503-681-9968 Fax: 503-844-7770

POST-OPERATIVE ORAL SEDATION INSTRUCTIONS

1. PATIENT SHOULD NOT DRIVE A MOTOR VEHICLE FOR 24 HOURS AFTER SEDATION
2. DO NOT OPERATE ANY HAZARDOUS DEVICES/MACHINERY FOR 24 HOURS AFTER SEDATION
3. A RESPONSIBLE ADULT PERSON SHOULD BE WITH THE PATIENT UNTIL HE/SHE HAS FULLY RECOVERED FROM THE EFFECTS OF THE SEDATION.
4. THE PATIENT SHOULD NOT GO UP AND DOWN STAIRS UNATTENDED. WHENEVER POSSIBLE, HAVE THE PATIENT STAY ON THE FIRST FLOOR UNTIL RECOVERED.
5. THE PATIENT SHOULD RESUME NORMAL EATING AND DRINKING AFTER THE SEDATION APPOINTMENT, UNLESS OTHERWISE INSTRUCTED BY THE DENTIST.
6. THE PATIENT NEEDS TO DRINK PLENTY OF WATER POST-OPERATIVELY TO PREVENT DEHYDRATION AND TO PREVENT THE "HANG-OVER" OF SEDATION MEDICATIONS.
7. AFTER LEAVING THE DENTAL OFFICE, PATIENTS SHOULD NOT BE LEFT UNATTENDED. PATIENTS MAY SEEM NORMAL AND RECOVERED, BUT THE EFFECTS OF THE MEDICATION CAN LAST FOR SEVERAL HOURS AFTER THE APPOINTMENT HAS ENDED. DO NOT ALLOW THE PATIENT TO MAKE IMPORTANT DECISIONS FOR THE REMAINDER OF THE DAY FOLLOWING HIS/HER SEDATION APPOINTMENT.
8. ALWAYS HOLD PATIENT'S ARM WHEN WALKING AS THEY MAY HAVE PROBLEMS WITH BALANCE WHILE UNDER THE EFFECTS OF THE SEDATIVE MEDICATIONS.
9. CALL THE OFFICE IF YOU HAVE ANY QUESTIONS OR CONCERNS. IF YOU FEEL THAT YOUR SYMPTOMS WARRANT A PHYSICIAN AND YOU ARE UNABLE TO REACH US, GO TO THE NEAREST EMERGENCY ROOM IMMEDIATELY.

FOLLOWING MOST SURGICAL PROCEDURES THERE MAY OR MAY NOT BE PAIN. YOU WILL BE PROVIDED WITH MEDICATION FOR DISCOMFORT THAT IS APPROPRIATE FOR YOU. IN MOST CASES, A NON-NARCOTIC PAIN REGIMEN IS RECOMMENDED CONSISTING OF *ACETAMINOPHEN* (TYLENOL®) AND *IBUPROFEN* (ADVIL®). THESE TWO MEDICATIONS **TAKEN TOGETHER**, CAN BE AS EFFECTIVE AS A NARCOTIC WITHOUT ANY OF THE SIDE AFFECTS. IF A NARCOTIC HAS BEEN PRESCRIBED, FOLLOW THE DIRECTIONS CAREFULLY. IF YOU HAVE ANY QUESTIONS ABOUT THESE MEDICATIONS INTERACTING WITH OTHER MEDICATIONS YOU ARE PRESENTLY TAKING, PLEASE CALL OUR OFFICE, YOUR PHYSICIAN AND/OR YOUR PHARMACIST.

Companion Signature: _____ TIME/DATE: _____